

NHSGGC Non haematology SACT regimes - guidance on which patients should receive seasonal influenza vaccine

Site Specific Team	Regime	Comments
Gyn/Sarcoma	CBOPBEP	OK before - not once started
Gyn/Sarcoma	Carboplatin	OK at any time - try to avoid 10-17 days post carboplatin
Gyn/Sarcoma	BEP	OK before - not once started
Gyn/Sarcoma	TIP	OK before - not once started
Gyn/Sarcoma	VIDE	OK before - not once started
Gyn/Sarcoma	VDC/IE	OK before - not once started
Gyn/Sarcoma	VAI	OK before - not once started
Gyn/Sarcoma	EUROBOSS	OK before - not once started
Gyn/Sarcoma	MAP	OK before - not once started
Gyn/Sarcoma	Mifurmatide	Not to be given
Gyn/Sarcoma	IVAdox	OK before - not once started
Gyn/Sarcoma	Doxorubicin	Leading up to next cycle
Gyn/Sarcoma	Imatinib	Any time
Gyn/Sarcoma	Sunitinib	Any time
Gyn/Sarcoma	Regorafenib	Any time
Gyn/Sarcoma	Pazopanib	Any time
Urology - Prostate	Docetaxel	OK before, during or after
Urology - Prostate	Abiraterone	(Note: technically SACT) Good idea before, during or after. Any time in cycle. These patients are on steroids
Urology - Prostate	Enzalutamide	(Note: technically SACT) but not necessarily on steroids. Not specifically indicated, but appropriate any time
Urology - Bladder	Carbo-gem	As per as per standard guidance, best given in 1st or 3rd week to avoid thrombocytopenia
Urology - Bladder	Cis-gem	As per as per standard guidance, best given in first or 3rd week to avoid thrombocytopenia
Urology - Bladder	Weekly paclitaxel	Any time
Urology - Renal	Sunitinib	Any time
Urology - Renal	Pazopanib	Any time
Urology - Renal	Axitinib	Any time
Urology - Renal	Everolimus	Any time, but ideally before starting
Breast	all regimes as per standard guidance	Vaccination should not delay neo-adjuvant chemotherapy
Gyn	all regimes as per standard guidance	Not to be given if on clinical trial with nivolumab/ ipilimumab
Melanoma	Pembrolizumab/ Ipilimumab/ Nivolumab/combination	Pre-treatment or just prior to next cycle. If on a clinical trial check protocol
Melanoma	Dabrafenib/ combination with tremetanib	Any time
CNS	all regimes as per standard guidance	Optimal timing during week 4 for Temozolomide 4 weekly cycles
GI	all regimes as per standard guidance	Timing as per guidance letter
Head and Neck	all regimes as per standard guidance	Timing as per guidance letter
Lung	All regimes except immunotherapy.	Timing as per guidance letter. If on a clinical trial check protocol
Lung	Immunotherapy: Nivolumab, pembrolizumab	Any time. If on a clinical trial check protocol.