

Evidence Briefing 7: Adults

Need to know

- For the purposes of this briefing, adults have been defined as anyone aged 25-50 years old.
- 24% of adults in Scotland exceeded the low-risk weekly drinking guidelines in 2017¹
- In 2014/15, 6% of people in Scotland had used one or more illicit drugs in the last year²
- Problem alcohol and drug use amongst adults are more prevalent in Greater Glasgow and Clyde than on average for Scotland³
- The rate of problem drug use amongst adults was highest in the 25 to 34 years age group in Scotland⁴
- On average, men consume alcohol on more days of the week than women in Scotland, and consume more units of alcohol⁵

Key findings

- There is strong evidence for the effectiveness of Alcohol Brief Interventions (ABIs) in primary care settings in reducing the weekly consumption of alcohol in adults⁶
- There is strong evidence for the effectiveness of well-planned psychosocial and developmental prevention interventions involving multiple services in reducing alcohol and drug related harms
- There is some evidence for the effectiveness of cognitive behavioural therapy, behavioural couples' therapy and pharmacotherapy in reducing alcohol and drug related harms, as well as clear alcohol and drug policies in the workplace
- There is an evidence gap relating to whether diversionary activities can be effective in preventing alcohol and drug use in adults

Good practice

- NHS Health Scotland resources on delivery of ABIs
- Oldham Borough Council pilots
- Brighton and Hove City Council 'named workers'
- Scotland's Take Home Naloxone Programme⁷
- Newcastle City Council roll-out of Naloxone
- Healthy Working Lives⁸

Potential Stakeholders

- Adult alcohol and drug services
- Allied health professionals
- Employers
- Scottish Prisons Service
- Recovery groups
- Local authority staff
- Social care staff
- Mental health professionals
- Community learning and development staff
- Police Scotland

1 Introduction

This is a briefing on what works for adults in alcohol and drug prevention. It relates to interventions for individuals between **the ages of 25 and 50**. However, it is important to note that there is a large crossover of what works with:

- The children and young people evidence briefing at the lower end of the 25 – 50 age range
- The older adults evidence briefing at the upper end of the 25 – 50 age range. In particular, individuals below 50 years of age may be presenting with alcohol and drug issues that may make the briefing on what works for older adults more appropriate.

There is evidence to suggest that sex and gender may affect how adults respond to alcohol and drugs, the risk and protective factors for reducing the likelihood of use, and the barriers to accessing support with this. Some studies have cited hormonal differences between the biological sexes as an important factor in use, recovery and relapse, with implications for what may work in intervention and treatment, particularly when pharmacological interventions are being considered⁹. Other differences are socio-cultural, and are thus more affected by societal structures¹⁰. The complexity of how gender and sex can affect alcohol and drug use and responses to treatment, as well as the overlap between gender and other risk factors (see At-risk briefing) make it difficult to determine how best to tailor alcohol and drug prevention programmes. However, it is clear that sex and gender should be taken into account in design and delivery of services.

2 Context

- 24% of adults in Scotland exceeded the low-risk weekly drinking guidelines in 2017¹¹
- In 2014/15, 6% of people in Scotland had used one or more illicit drugs in the last year¹²
- 22 people die every week in Scotland because of alcohol¹³
- There were 1,139 deaths in Scotland due to a cause wholly attributable to alcohol (alcohol-specific)¹⁴
- There were 1187 deaths from a drug overdose in 2018 in Scotland¹⁵
- Around 1.7 million working days are lost per year due to alcohol-related absence in Scotland¹⁶
- On average, men consume alcohol on more days of the week than women in Scotland, and consume more units of alcohol¹⁷
- There were 1,060 alcohol-related hospital stays¹⁸ and 238 drug-related hospital stays¹⁹ in Greater Glasgow and Clyde in 2016
- There are an estimated 18,700 people in Greater Glasgow and Clyde with problem drug use, almost three-quarters of whom are men²⁰
- One-third of people with problem drug use in Scotland are located in Greater Glasgow and Clyde²¹
- The rate of problem drug use amongst males and females was highest in the 25 to 34 years age group in Scotland²²

3 Detailed analysis of what works

Key to evidence briefing

High quality and multiple source evidence to support this approach (1)

Some evidence or emerging evidence to support this approach (2)

Limited evidence for this approach or potential development area for further investigation (3)

Primary Care	Workplace Based	Community Based
Brief interventions (1)	Brief interventions (1)	Brief interventions (1)
Integrated treatment programmes (1)	Integrated treatment programmes (1)	Integrated treatment programmes (1)
Pharmacotherapy (2)	Clear drug and alcohol policies (2)	Behavioural couples' therapy (2)
Behavioural couples' therapy (2)	Cognitive behavioural therapy (2)	Cognitive behavioural therapy (2)
Cognitive behavioural therapy (2)	Alcohol and drug testing with negative consequences (3)	Pharmacotherapy (2)
General counselling and psychodynamic therapies (3)	General counselling and psychodynamic therapies (3)	Diversionary activities (3)
Lived experience testimonials (3)	Self-help and mindfulness therapies (3)	Lived experience testimonials (3)
Self-help and mindfulness therapies (3)		General counselling and psychodynamic therapies (3)
		Self-help and mindfulness therapies (3)

Intervention Type	Description	Relevant group setting	Effectiveness	Factors that help (✓) or hinder (×) effectiveness	Good Practice
Alcohol Brief interventions (ABIs)	<p>Brief interventions are short, evidence based, structured conversations between a patient/client and a trained professional. They can last as little as 5-15 minutes, but duration should depend on situation, including time available, level of drinking involved and what the client wants.</p> <p>The conversation should primarily concern motivating and supporting the individual to think about and/or plan a change in their drinking behaviour (and potentially environment) in order to reduce their consumption and/or their risk of harm.</p>	Primary Care	<p>Effective</p> <p>Transferable (to certain settings)</p> <p>Sustained impact</p> <p>Works particularly well in primary care settings, such as when delivered by GPs and nurses. Some emerging evidence for effectiveness in criminal justice settings</p>	<ul style="list-style-type: none"> ✓ Non-confrontational²³ and person centred²⁴ ✓ Internationally validated assessment tools such as AUDIT-C or FAST²⁵ used to identify hazardous drinkers (where delivery is planned out with Primary Care, implementation should be tailored to the setting²⁶) ✓ Age-appropriate ✓ Proportionate to vulnerabilities²⁷ and person-centred²⁸ ✓ Used opportunistically as part of routine visit to primary care services²⁹ ✓ Delivered face-to-face³⁰ ✓ Delivery staff are well-trained in ABIs³¹ ✓ The patient is in control of their drinking and is helped to make changes for themselves³² ✓ Motivational interviewing and asset-based techniques are employed³³ ✓ When used in the workplace or outside primary care settings, embedding ABIs in wider health and wellbeing initiatives works well ✓ Dedicated treatment staffing and funding streams³⁴ 	<p>Alcohol Brief Interventions, Practitioner Training</p> <p>Alcohol Brief Interventions, Training for Trainers</p> <p>Delivery of Alcohol Brief Interventions resources, NHS Health Scotland³⁵</p>
		Workplace Based	<p>More studies in community settings would be useful – Health Scotland are currently reviewing ABIs in primary care and wider settings as part of the Alcohol Framework 2018</p>		
		Community Based	<ul style="list-style-type: none"> × Used with scare tactics and images³⁶ × Used in small workplaces where undertaking an ABI will affect employment × Requires separate visit³⁷ × Takes place more than 48 hours after initial visit³⁸ × There is no evidence to suggest that ABIs are effective in under 16s as this research has not been done³⁹ × Giving generic, unstructured advice 		

Intervention Type	Description	Relevant group setting	Effectiveness	Factors that help (✓) or hinder (×) effectiveness	Good Practice
Integrated treatment programmes for alcohol and drug use issues	Well-planned psychosocial and developmental prevention interventions involving multiple services	Primary Care	There are more studies to suggest effectiveness than ineffectiveness, but intervention type requires more robust evaluation	<ul style="list-style-type: none"> ✓ Integrated treatment programmes for women with problem alcohol and drug use appear to reduce use⁴⁰ ✓ Interventions are more cost-effective if embedded within wider, pre-existing programmes⁴¹ ✓ There is general evidence that holistic programmes are more effective than singular interventions ✓ Partners should include social care, community learning and development⁴², housing, homelessness and mental health services and prisons⁴³, and learning should be shared between these services⁴⁴ 	Oldham Borough Council pilots ⁴⁷ Brighton and Hove City Council 'named workers' ⁴⁸
		Workplace Based	Many studies are based in Sweden thus more research is needed on transferability	<ul style="list-style-type: none"> ✓ Stable housing and access to employment or training opportunities are important for sustaining recovery⁴⁵ ✓ Innovative programmes eg utilising technology⁴⁶ 	
		Community Based		<ul style="list-style-type: none"> × No evidence for inhibiting factors 	

Intervention Type	Description	Relevant group setting	Effectiveness	Factors that help (✓) or hinder (×) effectiveness	Good Practice
Cognitive behavioural therapy (CBT) for alcohol and drug issues	<p>“CBT is where the patient works collaboratively with a therapist to achieve specific treatment goals.</p> <p>These goals may include recognising the impact of behavioural and/or thinking patterns on feeling states and encouraging alternative cognitive and or behavioural coping skills and strategies to reduce the severity of target symptoms and problems.”⁴⁹</p>	Primary Care	Some evidence of effectiveness ⁵⁰ compared to traditional alcohol and drug treatments	<ul style="list-style-type: none"> ✓ Motivational interviewing approaches ⁵¹ ✓ Using contingency management (eg offering goods vouchers as an abstinence reward)⁵² ✓ Assessed for feasibility and acceptability in the workplace, including option of referral to external counselling services⁵³ ✓ In combination with clear alcohol and drug policies in the workplace, and as part of a wider wellbeing initiative eg stress management 	
		Workplace based		<ul style="list-style-type: none"> × There is mixed evidence for the effectiveness of CBT in combination with other treatments such as pharmacotherapy for alcohol and drug use disorders or contingency management⁵⁴ 	
		Community Based		<ul style="list-style-type: none"> × Confidentiality not guaranteed in workplace 	
Behavioural couples’ therapy (BCT)	<p>Couples-based therapy involves the spouse or partner “expressing active support” for the person with alcohol or drug issues, in reducing use.</p> <p>The therapy itself operates similarly to CBT but has a greater focus on “effective communication skills”⁵⁵. BCT requires sustained engagement from the couple, over several months⁵⁶.</p>	Primary Care	Some evidence of effectiveness ⁵⁷ compared to traditional drug and alcohol treatments, and to individual therapy ⁵⁸	<ul style="list-style-type: none"> ✓ BCT is more effective when only one partner has an alcohol or drug issue⁵⁹ ✓ BCT sessions should be structured and evidence-based⁶⁰ 	
		Community Based		No effect at two months but abstinence improvements at six-month follow	

Intervention Type	Description	Relevant group setting	Effectiveness	Factors that help (✓) or hinder (×) effectiveness	Good Practice
			up		
Pharmacotherapy	<p>Pharmacotherapy involves the use of prescription drugs to reduce alcohol and drug consumption and/or harm.</p> <p>Some, such as Disulfiram, are used as “aversion therapy”⁶², creating unpleasant reactions to the consumption of alcohol.</p> <p>Others, such as Naltrexone reduce the “desire for alcohol”⁶³ instead. Naloxone serves a different function, reducing the likelihood of opioid overdose by reversing the effects⁶⁴.</p>	Primary Care	Effect sizes are small, but better than control	<ul style="list-style-type: none"> ✓ Acamprosate, Naltrexone, Nalmefene and Disulfiram have been shown to have some effect in reducing alcohol harm⁶⁵ ✓ There is some evidence to suggest Acamprosate and Naltrexone are more cost-effective when used in combination⁶⁶ ✓ There is some evidence to suggest that Nalmefene and other pharmacotherapies work best when coupled with CBT, behavioural therapies or social network therapies⁶⁷ ✓ The provision of take-home Naloxone has been shown in small-scale studies to be an important part of preventing opioid overdose⁶⁸ ✓ Pharmacological interventions need to be based on NICE guidelines⁶⁹ 	<p>Newcastle City Council roll-out of Naloxone⁷⁰</p> <p>Barnsley Metropolitan Borough Council Naloxone pilot⁷¹</p> <p>Scottish Government Evaluation of Scotland's Take-Home Naloxone Programme⁷²</p>
		Community Based		<ul style="list-style-type: none"> × Lack of adequate training for staff and those with alcohol or drug use issues, particularly relating to Naloxone 	

Intervention Type	Description	Relevant group setting	Effectiveness	Factors that help (✓) or hinder (×) effectiveness	Good Practice
Clear alcohol and drug policies	Alcohol and drug policies set out guidelines, principles and procedures for approaching alcohol and drug use in the workplace. They may include training manuals with step-by-step guides on how to deal with different situations.	Workplace based		<ul style="list-style-type: none"> ✓ Embedded in wider health and wellbeing policies⁷³ ✓ Policies preventing discrimination in recruitment due to alcohol and drug use⁷⁴ ✓ Support for employers and employees on alcohol and drug use in the workplace⁷⁵ ✓ Ethos of improving workplace wellbeing⁷⁶ ✓ Guarantee confidentiality to all employees⁷⁷ ✓ Clearly communicated policies⁷⁸ ✓ Policies are developed with input from all stakeholders⁷⁹ 	The health and safety executive have provided a number of resources for employers ⁸⁰ Healthy Working Lives ⁸¹
				<ul style="list-style-type: none"> × Stigmatising policies⁸² 	
Diversionsary activities	<p>Diversionsary activities can refer to any activity designed to prevent alcohol and drug use or related harm, for example physical, leisure or social activities.</p> <p>At present there is a lack of</p>	Community based	There is very limited evidence on whether diversionsary activities can prevent alcohol and drug use or related harm	<ul style="list-style-type: none"> ✓ Develop new programmes and monitor for effectiveness 	

Intervention Type	Description	Relevant group setting	Effectiveness	Factors that help (✓) or hinder (×) effectiveness	Good Practice
	evidence as to whether diversionary activities can help prevent alcohol and drug use and related harm, and a concurrent need to develop and monitor innovative programmes with this specific aim.			× Insufficient evidence around enablers and inhibitors	
<p>Alcohol and drug testing with negative consequences</p> <p>General counselling and psychodynamic therapies</p> <p>Self-help and mindfulness therapies</p> <p>Lived experience testimonials</p>	Literature suggests that these four approaches are unlikely to reduce alcohol and drug use and related harms	N/A	There is very limited evidence that these approaches work in tackling alcohol and drug use and related harms	<p>× Mindfulness meditation specifically for reducing alcohol and drug use and related harm⁸³</p> <p>× Self-help based treatment specifically for reducing alcohol and drug use and related harm⁸⁴</p> <p>× General counselling and psychodynamic therapies specifically for reducing alcohol and drug use and related harm⁸⁵</p> <p>× Alcohol and drug testing with negative consequences and a negative focus⁸⁶</p> <p>× Using lived experience testimonials is associated with “no,</p>	N/A

Intervention Type	Description	Relevant group setting	Effectiveness	Factors that help (✓) or hinder (×) effectiveness	Good Practice
				or negative, preventative outcomes” ⁸⁷	

References

-
- ¹ Information available at: <https://www.gov.scot/publications/alcohol-framework-2018-preventing-harm-next-steps-changing-relationship-alcohol/>
- ² Information available at: <https://www.gov.scot/publications/scottish-crime-justice-survey-2014-15-drug-use/>
- ³ Information available at: <https://www.isdscotland.org/Health-Topics/Drugs-and-Alcohol-Misuse/Publications/2019-03-05/2019-03-05-Drug-Prevalence-2015-16-Report.pdf>
- ⁴ Ibid
- ⁵ Information available at: <https://www.gov.scot/binaries/content/documents/govscot/publications/statistics/2019/09/scottish-health-survey-2018-volume-1-main-report/documents/scottish-health-survey-2018-edition-volume-1-main-report/scottish-health-survey-2018-edition-volume-1-main-report/govscot%3Adocument/scottish-health-survey-2018-edition-volume-1-main-report.pdf>
- ⁶ *The Public Health Burden of Alcohol and the Effectiveness and Cost-Effectiveness of Alcohol Control Policies An evidence review*, Public Health England, 2016. Available at: https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/733108/alcohol_public_health_burden_evidence_review_updae_2018.pdf
- ⁷ National Naloxone Programme Scotland Monitoring Report 2017/18, ISD, <https://www.isdscotland.org/Health-Topics/Drugs-and-Alcohol-Misuse/Publications/2018-11-27/2018-11-27-Naloxone-Report.pdf>
- ⁸ Healthy Working Lives <https://www.healthyworkinglives.scot/Pages/default.aspx>
- ⁹ *Gender Considerations in Addiction: Implications for Treatment*, Polak, Haug, Drachenberg, Svikis, 2015. Available at: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4578628/>
- ¹⁰ *Sex differences, gender and addiction*, Becker, McClellan, Glover Reed, 2017. Available at: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5120656/>
- ¹¹ Information available at: <https://www.gov.scot/publications/alcohol-framework-2018-preventing-harm-next-steps-changing-relationship-alcohol/>
- ¹² Information available at: <https://www.gov.scot/publications/scottish-crime-justice-survey-2014-15-drug-use/>
- ¹³ Information available at: <http://www.healthscotland.scot/media/1863/mesas-monitoring-report-2018.pdf>
- ¹⁴ Information available at: <https://www.alcohol-focus-scotland.org.uk/alcohol-information/alcohol-facts-and-figures/>
- ¹⁵ Information available at: <https://www.nrscotland.gov.uk/statistics-and-data/statistics/statistics-by-theme/vital-events/deaths/drug-related-deaths-in-scotland/2018>
- ¹⁶ Information available at: <https://www.gov.scot/publications/scottish-health-survey-2017-volume-1-main-report/pages/51/>
- ¹⁷ Ibid
- ¹⁸ Information available at: <https://scotpho.nhsns.scot.nhs.uk/scotpho/spineChartAction.do>
- ¹⁹ Ibid
- ²⁰ Information available at: <https://www.isdscotland.org/Health-Topics/Drugs-and-Alcohol-Misuse/Publications/2019-03-05/2019-03-05-Drug-Prevalence-2015-16-Report.pdf>
- ²¹ Ibid
- ²² Ibid
- ²³ *Local Delivery Plan Standard: Alcohol Brief Interventions, National Guidance: 2019-20*, The Scottish Government. Available at: <https://www.gov.scot/publications/alcohol-brief-interventions-national-guidance/>
- ²⁴ *Local Delivery Plan Standard: Alcohol Brief Interventions, National Guidance: 2019-20*, The Scottish Government. Available at: <https://www.gov.scot/publications/alcohol-brief-interventions-national-guidance/>
- ²⁵ *Delivering Alcohol Brief Interventions*, NHS Health Scotland, 2017. Available at: http://www.healthscotland.scot/media/1361/abi-briefing-paper_apr2017_english.pdf
- ²⁶ *Impact of alcohol: exploring the learning from elsewhere*, Kincaid, Montgomery, 2018. Available at: https://effectiveservices.org/downloads/Impact_of_Alcohol_Three-02-08-2018.pdf?platform=hootsuite
- ²⁷ *Drug misuse prevention: targeted interventions*, NICE guideline, Public Health England, February 2017. Available at: <https://www.nice.org.uk/guidance/ng64>
- ²⁸ *Local Delivery Plan Standard: Alcohol Brief Interventions, National Guidance: 2019-20*, The Scottish Government. Available at: <https://www.gov.scot/publications/alcohol-brief-interventions-national-guidance/>

²⁹ Ibid

³⁰ Ibid

³¹ Ibid

³² *Delivering Alcohol Brief Interventions*, NHS Health Scotland, 2017. Available at: http://www.healthscotland.scot/media/1361/abi-briefing-paper_apr2017_english.pdf

³³ *Local Delivery Plan Standard: Alcohol Brief Interventions, National Guidance: 2019-20*, The Scottish Government. Available at: <https://www.gov.scot/publications/alcohol-brief-interventions-national-guidance/>

³⁴ *The Public Health Burden of Alcohol and the Effectiveness and Cost-Effectiveness of Alcohol Control Policies An evidence review*, Public Health England, 2016. Available at: https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/733108/alcohol_public_health_burden_evidence_review_update_2018.pdf

³⁵ More information available at: <http://www.healthscotland.scot/publications/alcohol-brief-intervention-resources>

³⁶ *The international evidence on the prevention of drug and alcohol use Summary and examples of implementation in England*, Public Health England, 2015. Available at: https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/669654/Preventing_drug_and_alcohol_misuse_international_evidence_and_implementation_examples.pdf

³⁷ *Local Delivery Plan Standard: Alcohol Brief Interventions, National Guidance: 2019-20*, The Scottish Government. Available at: <https://www.gov.scot/publications/alcohol-brief-interventions-national-guidance/>

³⁸ Ibid

³⁹ *Local Delivery Plan Standard: Alcohol Brief Interventions, National Guidance: 2019-20*, The Scottish Government. Available at: <https://www.gov.scot/publications/alcohol-brief-interventions-national-guidance/>

⁴⁰ *Evidence summary: public health interventions in response to substance misuse (drugs) to support parents, their children and young people up to 25 years*, NHS Health Scotland, 2014. Available at: <http://www.healthscotland.com/documents/24075.aspx>

⁴¹ *The Public Health Burden of Alcohol and the Effectiveness and Cost-Effectiveness of Alcohol Control Policies An evidence review*, Public Health England, 2016. Available at: https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/733108/alcohol_public_health_burden_evidence_review_update_2018.pdf

⁴² *Review of Community Alcohol Campaigns Recommendations*, Dudleston Harkins Social Research Limited, 2014. Available at: <https://www.glasgow.gov.uk/councillorsandcommittees/viewSelectedDocument.asp?c=P62AFQDNT12UT1T1DN>

⁴³ *Preventing Drug-related Deaths*, Local Government Association, 2017. Available at: <https://www.local.gov.uk/preventing-drug-related-deaths>

⁴⁴ *Coexisting severe mental illness and substance misuse: community health and social care services*, NICE, 2016. Available at: <https://www.nice.org.uk/guidance/ng58>

⁴⁵ *An independent review into the impact on employment outcomes of drug or alcohol addiction, and obesity*, Black, 2016. Available at: https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/573892/employment-outcomes-of-drug-or-alcohol-addiction-and-obesity-print.PDF

⁴⁶ *Local Solutions, Healthy Lives: council's role in drug and alcohol services*, Local Government Association, 2015. Available at: <https://www.bl.uk/britishlibrary/~media/bl/global/social-welfare/pdfs/non-secure/l/o/c/local-solutions--healthy-lives-councils-role-in-drug-and-alcohol-services.pdf>

⁴⁷ Ibid

⁴⁸ Ibid

⁴⁹ *The Public Health Burden of Alcohol and the Effectiveness and Cost-Effectiveness of Alcohol Control Policies An evidence review*, Public Health England, 2016. Available at: https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/733108/alcohol_public_health_burden_evidence_review_update_2018.pdf

⁵⁰ *The Public Health Burden of Alcohol and the Effectiveness and Cost-Effectiveness of Alcohol Control Policies An evidence review*, Public Health England, 2016. Available at: https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/733108/alcohol_public_health_burden_evidence_review_update_2018.pdf

⁵¹ *Cognitive-Behavioural Therapy for Substance Use Disorders*, McHugh, Hearon, Otto, 2010. Available at: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2897895>

⁵² Ibid

⁵³ *Counselling in the Workplace: a comprehensive review of the research- 2nd edition*, British Association for Counselling and Psychotherapy, 2008. Available at: <https://www.bacp.co.uk/media/1974/bacp-counselling-in-workplace-systematic-review.pdf>

⁵⁴ *Cognitive-Behavioural Therapy for Substance Use Disorders*, McHugh, Hearon, Otto, 2010. Available at: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2897895>

-
- ⁵⁵ *The Public Health Burden of Alcohol and the Effectiveness and Cost-Effectiveness of Alcohol Control Policies An evidence review*, Public Health England, 2016. Available at: https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/733108/alcohol_public_health_burden_evidence_review_update_2018.pdf
- ⁵⁶ *Behavioural Couples Therapy for Substance Abuse: Rationale, Methods, and Findings*, Fals-Stewart, O'Farrell, Birchler, 2004. Available at: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2851021/>
- ⁵⁷ *The Public Health Burden of Alcohol and the Effectiveness and Cost-Effectiveness of Alcohol Control Policies An evidence review*, Public Health England, 2016. Available at: https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/733108/alcohol_public_health_burden_evidence_review_update_2018.pdf
- ⁵⁸ *Behavioural Couples Therapy for Substance Abuse: Rationale, Methods, and Findings*, Fals-Stewart, O'Farrell, Birchler, 2004. Available at: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2851021/>
- ⁵⁹ Ibid
- ⁶⁰ Ibid
- ⁶¹ Ibid
- ⁶² *The Public Health Burden of Alcohol and the Effectiveness and Cost-Effectiveness of Alcohol Control Policies An evidence review*, Public Health England, 2016. Available at: https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/733108/alcohol_public_health_burden_evidence_review_update_2018.pdf
- ⁶³ Ibid
- ⁶⁴ *Preventing Drug-related Deaths: case studies*, Local Government Association, 2017. Available at: <http://ow.ly/Gw0430ehjsH>
- ⁶⁵ *The Public Health Burden of Alcohol and the Effectiveness and Cost-Effectiveness of Alcohol Control Policies An evidence review*, Public Health England, 2016. Available at: https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/733108/alcohol_public_health_burden_evidence_review_update_2018.pdf
- ⁶⁶ Ibid
- ⁶⁷ Ibid
- ⁶⁸ *Preventing Drug-related Deaths*, Local Government Association, 2017. Available at: <https://www.local.gov.uk/preventing-drug-related-deaths>
- ⁶⁹ Guidance available at: <https://www.nice.org.uk/guidance/published?type=apg,csg,cg,mpg,ph,sg,sc>
- ⁷⁰ *Preventing Drug-related Deaths*, Local Government Association, 2017. Available at: https://www.local.gov.uk/sites/default/files/documents/22.3%20Drug%20related%20deaths_v8.pdf
- ⁷¹ Ibid
- ⁷² National Naloxone Programme Scotland Monitoring Report 2017/18, ISD, <https://www.isdscotland.org/Health-Topics/Drugs-and-Alcohol-Misuse/Publications/2018-11-27/2018-11-27-Naloxone-Report.pdf>
- ⁷³ Resource available at: https://www.ilo.org/safework/info/instr/WCMS_178438/lang--en/index.htm
- ⁷⁴ *International Standards on Drug Use Prevention, Second Updated Edition*, World Health Organisation/United Nations Office on Drugs and Crime, 2018. Available at: <http://www.unodc.org/unodc/en/prevention/workplaceprevention.html>
- ⁷⁵ Ibid
- ⁷⁶ *Easing the transition: The relationship between alcohol and labour market activity in the over 50s population of the UK*, Drink Wise Age Well, 2016. Available at: <https://ilcuk.org.uk/easing-the-transition-the-relationship-between-alcohol-and-labour-market-activity-in-the-over-50s-population-of-the-uk/>
- ⁷⁷ *The international evidence on the prevention of drug and alcohol use Summary and examples of implementation in England*, Public Health England, 2015. Available at: https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/669654/Preventing_drug_and_alcohol_misuse_international_evidence_and_implementation_examples.pdf
- ⁷⁸ *International Standards on Drug Use Prevention, Second Updated Edition*, World Health Organisation/United Nations Office on Drugs and Crime, 2018. Available at: <http://www.unodc.org/unodc/en/prevention/prevention-standards.html>
- ⁷⁹ Ibid
- ⁸⁰ Resources available at: <http://www.hse.gov.uk/alcoholdrugs/>
- ⁸¹ Healthy Working Lives <https://www.healthyworkinglives.scot/Pages/default.aspx>

⁸² *International Standards on Drug Use Prevention, Second Updated Edition*, World Health Organisation/United Nations Office on Drugs and Crime, 2018. Available at:

<http://www.unodc.org/unodc/en/prevention/prevention-standards.html>

⁸³ *The Public Health Burden of Alcohol and the Effectiveness and Cost-Effectiveness of Alcohol Control Policies An evidence review*, Public Health England, 2016. Available at:

https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/733108/alcohol_public_health_burden_evidence_review_update_2018.pdf

⁸⁴ Ibid

⁸⁵ Ibid

⁸⁶ *The international evidence on the prevention of drug and alcohol use Summary and examples of implementation in England*, Public Health England, 2015. Available at:

https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/669654/Preventing_drug_and_alcohol_misuse_international_evidence_and_implementation_examples.pdf

⁸⁷ *Alcohol Framework 2018: Preventing Harm next steps on changing our relationship with alcohol*, Scottish Government, 2018. Available at: <https://www.gov.scot/publications/alcohol-framework-2018-preventing-harm-next-steps-changing-relationship-alcohol/>